

## References

1. Physical Activity Guidelines Advisory Committee I. 2018. Physical Activity Guidelines Advisory Committee scientific report. Washington: U.S. Department of Health and Human Services; 2018
2. Bø K. Urinary incontinence, pelvic floor dysfunction, exercise and sport. *Sports Med.* 2004;34(7):451–64.
3. Nygaard IE, Shaw JM. Physical activity and the pelvic floor. *Am J Obstet Gynecol.* 2016;214(2):164–71.
4. Constantinou CE, Govan DE. Spatial distribution and timing of transmitted and reflexly generated urethral pressures in healthy women. *J Urol.* 1982;127(5):964–9.
5. Miller JM, Ashton-Miller JA, DeLancey JO. A pelvic muscle precontraction can reduce cough-related urine loss in selected women with mild SUI. *J Am Geriatr Soc.* 1998;46(7):870–4.
6. Seegmiller JG, McCaw ST. Ground reaction forces among gymnasts and recreational athletes in drop landings. *J Athl Train.* 2003;38(4):311–4.
7. James ED. The behaviour of the bladder during physical activity. *Br J Urol.* 1978;50(6):387–94.
8. Ree ML, Nygaard I, Bø K. Muscular fatigue in the pelvic floor muscles after strenuous physical activity. *Acta Obstet Gynecol Scand.* 2007;86(7):870–6.
9. Middlekauff ML, Egger MJ, Nygaard IE, Shaw JM. The impact of acute and chronic strenuous exercise on pelvic floor muscle strength and support in nulliparous healthy women. *Am J Obstet Gynecol.* 2016;215(3):316.e1–7.
10. de Araujo M, Parmigiano T, Della Negra L, Torelli L, de Carvalho C, Wo L, et al. Evaluation of athletes' pelvic floor: is there a relation with urinary incontinence? *Rev Bras Med Esporte.* 2015;21:442–6.
11. Jürgensen SP, Borghi-Silva A, Bastos AMFG, Correia GN, Pereira-Baldon VS, Cabiddu R, et al. Relationship between aerobic capacity and pelvic floor muscles function: a cross-sectional study. *Braz J Med Biol Res.* 2017;50(11):e5996
12. Bø K, Ellstrøm Engh M, Hilde G. Regular exercisers have stronger pelvic floor muscles than nonregular exercisers at midpregnancy. *Am J Obstet Gynecol.* 2018;218(4):427.e1–5.
13. Varella LR, Torres VB, Angelo PH, Eugênia de Oliveira MC, Matias de Barros AC, Viana EeS, et al. Influence of parity, type of delivery, and physical activity level on pelvic floor muscles in postmenopausal women. *J Phys Ther Sci.* 2016;28(3):824–30.
14. Ferla L, Paiva LL, Darki C, Vieira A. Comparison of the functionality of pelvic floor muscles in women who practice the Pilates method and sedentary women: a pilot study. *Int Urogynecol J.* 2016;27(1):123–8.

15. Moss W, Shaw JM, Yang M, Sheng X, Hitchcock R, Niederauer S, et al. The association between pelvic floor muscle force and general strength and fitness in postpartum women. *Female Pelvic Med Reconstr Surg*. 2019. <https://doi.org/10.1097/SPV.0000000000000718>.
16. Carvalhais A, Da Roza T, Vilela S, Jorge RN, Bø K. Association between physical activity level and pelvic floor muscle variables in women. *Int J Sports Med*. 2018;39(13):995–1000.
17. Luginbuehl H, Naeff R, Zahnd A, Baeyens JP, Kuhn A, Radlinger L. Pelvic floor muscle electromyography during different running speeds: an exploratory and reliability study. *Arch Gynecol Obstet*. 2016;293(1):117–24.
18. Saeuberli PW, Schraknepper A, Eichelberger P, Luginbuehl H, Radlinger L. Reflex activity of pelvic floor muscles during drop landings and mini-trampolining-exploratory study. *Int Urogynecol J*. 2018;29(12):1833–40.
19. Moser H, Leitner M, Baeyens JP, Radlinger L. Pelvic floor muscle activity during impact activities in continent and incontinent women: a systematic review. *Int Urogynecol J*. 2018;29(2):179–96.
20. Kruger JA, Murphy BA, Heap SW. Alterations in levator ani morphology in elite nulliparous athletes: a pilot study. *Aust N Z J Obstet Gynaecol*. 2005;45(1):42–7.
21. Kruger JA, Dietz HP, Murphy BA. Pelvic floor function in elite nulliparous athletes. *Ultrasound Obstet Gynecol*. 2007;30(1):81–5.
22. Roza TD, Brandão S, Oliveira D, Mascarenhas T, Parente M, Duarte JA, et al. Football practice and urinary incontinence: relation between morphology, function and biomechanics. *J Biomech*. 2015;48(9):1587–92.
23. Hannestad YS, Rortveit G, Daltveit AK, Hunskaar S. Are smoking and other lifestyle factors associated with female urinary incontinence? The Norwegian EPINCONT Study. *BJOG*. 2003;110(3):247–54.
24. [Janeisa Franck Virtuoso](#)<sup>1</sup>, [Enaiane Cristina Menezes](#)<sup>2</sup>, [Giovana Zarpellon Mazo](#)<sup>2</sup> (2019) Effect of Weight Training with Pelvic Floor Muscle Training in Elderly Women with Urinary Incontinence June ;90(2):141-150 Epub April 4